

(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class – XI)

Miscellaneous Exercise on chapter 1

Question 1:

Decide, among the following sets, which sets are subsets of one and another: $A = \{x: x \in \mathbb{R} \text{ and } x \text{ satisfy } x^2 - 8x + 12 = 0\}, B = \{2, 4, 6\},$ $C = \{2, 4, 6, 8...\}, D = \{6\}.$

Answer 1:

A = { $x: x \in R$ and x satisfies $x^2 - 8x + 12 = 0$ } 2 and 6 are the only solutions of $x^2 - 8x + 12 = 0$. $\therefore A = \{2, 6\}$ B = {2, 4, 6}, C = {2, 4, 6, 8 ...}, D = {6} $\therefore D \subset A \subset B \subset C$ Hence, $A \subset B$, $A \subset C$, $B \subset C$, $D \subset A$, $D \subset B$, $D \subset C$

Question 2:

In each of the following, determine whether the statement is true or false. If it is true, prove it. If it is false, give an example.

- (i) If $x \in A$ and $A \in B$, then $x \in B$
- (ii) If $A \subset B$ and $B \in C$, then $A \in C$
- (iii) If $A \subset B$ and $B \subset C$, then $A \subset C$
- (iv) If $A \not\subset B$ and $B \not\subset C$, then $A \not\subset C$
- (v) If $x \in A$ and $A \not\subset B$, then $x \in B$
- (vi) If $A \subset B$ and $x \notin B$, then $x \notin A$

Answer 2:

(i) False Let A = {1, 2} and B = {1, {1, 2}, {3}}



(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class – XI)

Now,

 $2 \in \{1, 2\}$ and $\{1, 2\} \in \{\{3\}, 1, \{1, 2\}\}$

$\because \mathsf{A} \in \mathsf{B}$

However,

 $2 \notin \{\{3\}, 1, \{1, 2\}\}$

(ii) False

Let

 $A = \{2\}, B = \{0, 2\}, and C = \{1, \{0, 2\}, 3\}$ As $A \subset B$ $B \in C$ However, $A \notin C$ (iii) True
Let $A \subset B$ and $B \subset C$.
Let $x \in A$ $\Rightarrow x \in B \qquad [\because A \subset B]$ $\Rightarrow x \in C \qquad [\because B \subset C]$ $\therefore A \subset C$ (iv) False $A = \{1, 2\}, B = \{0, 6, 8\}, and C = \{0, 1, 2, 6, 9\}$ $A \not\subset B \text{ and } B \not\subset C$ Let
Accordingly,

However, $A \subset C$ (v) False Let $A = \{3, 5, 7\}$ and $B = \{3, 4, 6\}$ Now, $5 \in A$ and $A \notin B$ However, $5 \notin B$



Eiwar

(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class - XI)


(vi) True Let $A \subset B$ and $x \notin B$. To show: $x \notin A$ If possible, suppose $x \in A$. Then, $x \in B$, which is a contradiction as $x \notin B$ $\therefore x \notin A$

Question 3:

Let A, B and C be the sets such that A \cup B = A \cup C and A \cap B = A \cap C. show that B = C.

Answer 3:

Let, A, B and C be the sets such that $A \cup B = A \cup C$ and $A \cap B = A \cap C$. To show: B = CLet $x \in B$ $[B \subset A \cup B]$ $\Rightarrow x \in A \cup B$ $[A \cup B = A \cup C]$ $\Rightarrow x \in A \cup C$ $\Rightarrow x \in A \text{ or } x \in C$ Case I $x \in$ А Also, $x \in B$ $x \in A \cap B$ ÷ $\Rightarrow x \in A \cap C \quad [:: A \cap B = A \cap C]$ $\therefore x \in A \text{ and } x \in C$ $\therefore x \in C$ $\therefore B \subset C$ Similarly, we can show that $C \subset B$. \therefore B = C



(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class – XI)

Question 4:

Show that the following four conditions are equivalent:

(i) $A \subset B$ (ii) $A - B = \Phi$ (iii) $A \cup B = B$ (iv) $A \cap B = A$

Answer 4:

First, we have to show that (i) \Leftrightarrow (ii). Let $A \subset B$ To show: A – B $\neq \Phi$ If possible, suppose A – B $\neq \Phi$ This means that there exists $x \in A$, $x \neq B$, which is not possible as $A \subset B$. $\therefore A - B = \Phi$ $\therefore A \subset B \Rightarrow A - B = \Phi$ Let A – B = Φ To show: $A \subset B$ Let $x \in A$ Clearly, $x \in B$ because if $x \notin B$, then $A - B \neq \Phi$ $\therefore A - B = \Phi \Rightarrow A \subset B$ \therefore (i) \Leftrightarrow (ii) Let $A \subset B$ To show: $A \cup B = B$ Clearly, $B \subset A \cup B$ Let $x \in A \cup B$ $\Rightarrow x \in A \text{ or } x \in B$ Case I: $x \in A$ $[:: A \subset B]$ $\Rightarrow x \in \mathbf{B}$ $A \cup B \subset B$ Case II: $x \in B$ Then, $A \cup B = B$



Eiwar



(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class – XI)

Conversely, let $A \cup B = B$ Let $x \in A$ $[:: A \subset A \cup B]$ $\Rightarrow x \in A \cup B$ $[:: A \cup B = B]$ $\Rightarrow x \in \mathbf{B}$ $\therefore A \subset B$ Hence, (i) \Leftrightarrow (iii) Now, we have to show that (i) \Leftrightarrow (iv). Let $A \subset B$ $Clearly A \cap B \subset A$ Let $x \in A$ We have to show that $x \in A \cap B$ As $A \subset B$, $x \in B$ $x \in \mathbb{A} \cap \mathbb{B}$:. $A \subset A \cap B$ Hence, $A = A \cap B$ Conversely, suppose $A \cap B = A$ Let $x \in A$ $\Rightarrow x \in A \cap B$ \Rightarrow *x* \in A and *x* \in B $\Rightarrow x \in B$ $\therefore A \subset B$ Hence, (i) \Leftrightarrow (iv).

Question 5:

Show that if $A \subset B$, then $C - B \subset C - A$.

Answer 5:

Let $A \subset B$



Eiwar

(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class - XI)

To show: $C - B \subset C - A$ Let $x \in C - B$ $\Rightarrow x \in C$ and $x \notin B$ $\Rightarrow x \in C$ and $x \notin A$ [$A \subset B$] $\Rightarrow x \in C - A$ $\therefore C - B \subset C - A$

Question 6:

Assume that P(A) = P(B). Show that A = B.

Answer 6:

Let P(A) = P(B)To show: A = BLet $x \in A$ $A \in P(A) = P(B)$ $\therefore x \in C$, for some $C \in P(B)$ Now, $C \subset B$ $\therefore x \in B$ $\therefore A \subset B$ Similarly, $B \subset A$ $\therefore A = B$

Question 7:

Is it true that for any sets A and B, P (A) \cup P (B) = P (A \cup B)? Justify your answer.

Answer 7:

False Let A = $\{0, 1\}$ and B = $\{1, 2\}$

Eiwa

(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class – XI)

 $\therefore A \cup B = \{0, 1, 2\}$ $P(A) = \{\Phi, \{0\}, \{1\}, \{0, 1\}\}$ $P(B) = \{\Phi, \{1\}, \{2\}, \{1, 2\}\}$ $P(A \cup B) = \{\Phi, \{0\}, \{1\}, \{2\}, \{0, 1\}, \{1, 2\}, \{0, 2\}, \{0, 1, 2\}\}$ $P(A) \cup P(B) = \{\Phi, \{0\}, \{1\}, \{0, 1\}, \{2\}, \{1, 2\}\}$ $\therefore P(A) \cup P(B) \neq P(A \cup B)$

Question 8:

Show that for any sets A and B, A = $(A \cap B) \cup (A - B)$ and A $\cup (B - A) = (A \cup B)$

Answer 8:

To show: $A = (A \cap B) \cup (A - B)$ Let $x \in A$ We have to show that $x \in (A \cap B) \cup (A$ – B) Case I $x \in A \cap B$ Then, $x \in (A \cap B) \subset (A \cup B) \cup (A -$ B) Case II $x \notin A \cap B$ $\Rightarrow x \notin A \text{ or } x \notin B$ $\therefore x \notin B [x \notin A]$ $\therefore x \notin A - B \subset (A \cup B) \cup (A - B)$ $\therefore A \subset (A \cap B) \cup (A - B) \dots (1)$ It is clear that $A \cap B \subset A$ and $(A - B) \subset A$ $\therefore (A \cap B) \cup (A - B) \subset A \dots (2)$ From (1) and (2), we obtain $A = (A \cap B) \cup (A - B)$ To prove: $A \cup (B - A) \subset A \cup B$ Let $x \in A \cup (B - A)$



(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class - XI)

8

 $\Rightarrow x \in A \text{ or } x \in (B - A)$ $\Rightarrow x \in A \text{ or } (x \in B \text{ and } x \notin A)$ $\Rightarrow (x \in A \text{ or } x \in B) \text{ and } (x \in A \text{ or } x \notin A)$ $\Rightarrow x \in (A \cup B)$ $\therefore A \cup (B - A) \subset (A \cup B) \dots (3)$ Next, we show that $(A \cup B) \subset A \cup (B - A)$. Let $y \in A \cup B$ $\Rightarrow y \in A \text{ or } y \in B$ $\Rightarrow (y \in A \text{ or } y \in B) \text{ and } (y \in A \text{ or } y \notin A)$ $\Rightarrow y \in A \text{ or } (y \in B \text{ and } y \notin A)$ $\Rightarrow y \in A \cup (B - A)$ $\therefore A \cup B \subset A \cup (B - A) \dots (4)$ Hence, from (3) and (4), we obtain $A \cup (B - A) = A \cup B$.

Question 9:

Using properties of sets show that (i) $A \cup (A \cap B) = A$ (ii) $A \cap (A \cup B) = A$.

Answer 9:

(i) To show: $A \cup (A \cap B) = A$ We know that $A \subset A$ $A \cap B \subset A$ $\therefore A \cup (A \cap B) \subset A \dots (1)$ Also, $A \subset A \cup (A \cap B) \dots (2)$ \therefore From (1) and (2), $A \cup (A \cap B) = A$ (ii) To show: $A \cap (A \cup B) = A$ $A \cap (A \cup B) = (A \cap A) \cup (A \cap B)$ $= A \cup (A \cap B)$



(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class - XI)

= A {from (1)}

Question 10:

Show that $A \cap B = A \cap C$ need not imply B = C.

Answer 10:

Let A = $\{0, 1\}$, B = $\{0, 2, 3\}$, and C = $\{0, 4, 5\}$ Accordingly, A \cap B = $\{0\}$ and A \cap C = $\{0\}$ Here, A \cap B = A \cap C = $\{0\}$ However, B \neq C [2 \in B and 2 \notin C]

Question 11:


Let A and B be sets. If $A \cap X = B \cap X = \Phi$ and $A \cup X = B \cup X$ for some set X, show that A = B.

(Hints A = A \cap (A \cup X), B = B \cap (B \cup X) and use distributive law)

Answer 11:

Let A and B be two sets such that A \cap X = B \cap X = f and A \cup X = B \cup X for some set X.

To show: A = BIt can be seen that $A = A \cap (A \cup X) = A \cap (B \cup X) [A \cup X = B \cup X]$ $= (A \cap B) \cup (A \cap X) [Distributive law]$ $= (A \cap B) \cup \Phi [A \cap X = \Phi]$ $= A \cap B$(1) Now, $B = B \cap (B \cup X)$ $= B \cap (A \cup X) [A \cup X = B \cup X]$ $= (B \cap A) \cup (B \cap X) [Distributive law]$ $= (B \cap A) \cup \Phi [B \cap X = \Phi]$



(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class - XI)

= $B \cap A$ = $A \cap B$ (2) Hence, from (1) and (2), we obtain A = B.

Question 12:

Find sets A, B and C such that A \cap B, B \cap C and A \cap C are non-empty sets and A \cap B \cap C = Φ .

Answer 12:

Let A = {0, 1}, B = {1, 2}, and C = {2, 0}. Accordingly, A \cap B = {1}, B \cap C = {2}, and A \cap C = {0}. \therefore A \cap B, B \cap C, and A \cap C are non-empty. However, A \cap B \cap C = Φ

Question 13:

In a survey of 600 students in a school, 150 students were found to be taking tea and 225 taking coffee, 100 were taking both tea and coffee. Find how many students were taking neither tea nor coffee?

10

Answer 13:

Let U be the set of all students who took part in the survey. Let T be the set of students taking tea. Let C be the set of students taking coffee. Accordingly, n(U) = 600, n(T) = 150, n(C) = 225, $n(T \cap C) = 100$ To find: Number of student taking neither tea nor coffee i.e., we have to find $n(T' \cap C')$. $n(T' \cap C') = n(T \cup C)'$ $= n(U) - n(T \cup C)$ $= n(U) - [n(T) + n(C) - n(T \cap C)]$

(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class - XI)

= 600 - [150 + 225 - 100]

- = 600 275
- = 325

Hence, 325 students were taking neither tea nor coffee.

Question 14:

In a group of students 100 students know Hindi, 50 know English and 25 know both. Each of the students knows either Hindi or English. How many students are there in the group?

Answer 14:

Let U be the set of all students in the group. Let E be the set of all students who know English. Let H be the set of all students who know Hindi. \therefore H \cup E = U Accordingly, n(H) = 100 and n(E) = 50 $n(H \cup E) = n(H) + n(E) - n(H \cap E)$ = 100 + 50 - 25 = 125Hence, there are 125 students in the group.

Question 15:

In a survey of 60 people, it was found that 25 people read newspaper H, 26 read newspaper T, 26 read newspaper I, 9 read both H and I,11 read both H and T, 8 read both T and I, 3 read all three newspapers. Find:

(i) the number of people who read at least one of the newspapers.

(ii) the number of people who read exactly one newspaper.



(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class - XI)

Answer 15:

Let A be the set of people who read newspaper H. Let B be the set of people who read newspaper T. Let C be the set of people who read newspaper I. Accordingly,

 $n(A) = 25, n(B) = 26, n(C) = 26, n(A \cap C) = 9, n(A \cap B) = 11,$ $n(B \cap C) = 8, n(A \cap B \cap C) = 3$


Let U be the set of people who took part in the survey.

(i) Accordingly,

 $n(A \cup B \cup C) = n(A) + n(B) + n(C) - n(A \cap B) - n(B \cap C) - n(C \cap A) + n(A \cap B \cap C) = 25 + 26 + 26 - 11 - 8 - 9 + 3 = 52$

Hence, 52 people read at least one of the newspapers.

(ii) Let *a* be the number of people who read newspapers H and T only.



Let *b* denote the number of people who read newspapers I and H only. Let *c* denote the number of people who read newspapers T and I only. Let *d* denote the number of people who read all three newspapers. Accordingly, $d = n(A \cap B \cap C) = 3$

Now,
$$n(A \cap B) = a + d$$

$$n(B \cap C) = c + d$$
$$n(C \cap A) = b + d$$

 $\therefore a + d + c + d + b + d = 11 + 8 + 9 = 28$



(www.tiwariacademy.com : A step towards free education) (Chapter - 1) (Sets)

(Class – XI)

 $\Rightarrow a + b + c + d = 28 - 2d = 28 - 6 = 22$ Hence, (52 - 22) = 30 people read exactly one newspaper.

Question 16:


In a survey it was found that 21 people liked product A, 26 liked product B and 29 liked product C. If 14 people liked products A and B, 12 people liked products C and A, 14 people liked products B and C and 8 liked all the three products. Find how many liked product C only.

Answer 16:

Let A, B, and C be the set of people who like product A, product B, and product C respectively.

Accordingly, n(A) = 21, n(B) = 26, n(C) = 29, $n(A \cap B) = 14$, $n(C \cap A) = 12$, $n(B \cap C) = 14$, $n(A \cap B \cap C) = 8$

The Venn diagram for the given problem can be drawn as



It can be seen that number of people who like product C only is $\{29 - (4 + 8 + 6)\} = 11$

