

Mathematics

(www.tiwariacademy.com : A step towards free education)

(Chapter – 2) (Relations and Functions)

(Class – XI)

Miscellaneous Exercise on Chapter 2

Question 1:

The relation f is defined by $f(x) = \begin{cases} x^2, & 0 \leq x \leq 3 \\ 3x, & 3 \leq x \leq 10 \end{cases}$

The relation g is defined by $g(x) = \begin{cases} x^2, & 0 \leq x \leq 2 \\ 3x, & 2 \leq x \leq 10 \end{cases}$

Show that f is a function and g is not a function.

Answer 1:

The relation f is defined as

$$f(x) = \begin{cases} x^2, & 0 \leq x \leq 3 \\ 3x, & 3 \leq x \leq 10 \end{cases}$$

It is observed that for

$$0 \leq x < 3, \quad f(x) = x^2$$

$$3 < x \leq 10, \quad f(x) = 3x$$


Also, at $x = 3$, $f(x) = 3^2 = 9$ or $f(x) = 3 \times 3 = 9$ i.e., at $x = 3$, $f(x) = 9$

Therefore, for $0 \leq x \leq 10$, the images of $f(x)$ are unique. Thus, the given relation is a function.

The relation g is defined as

$$g(x) = \begin{cases} x^2, & 0 \leq x \leq 2 \\ 3x, & 2 \leq x \leq 10 \end{cases}$$

It can be observed that for $x = 2$, $g(x) = 2^2 = 4$ and $g(x) = 3 \times 2 = 6$



Mathematics

(www.tiwariacademy.com : A step towards free education)

(Chapter – 2) (Relations and Functions)

(Class – XI)

Hence, element 2 of the domain of the relation g corresponds to two different images i.e., 4 and 6.

Hence, this relation is not a function.

Question 2:

If $f(x) = x^2$, find. $\frac{f(1.1) - f(1)}{(1.1 - 1)}$

Answer 2:

$$f(x) = x^2$$

$$\therefore \frac{f(1.1) - f(1)}{(1.1 - 1)} = \frac{(1.1)^2 - (1)^2}{(1.1 - 1)} = \frac{1.21 - 1}{0.1} = \frac{0.21}{0.1} = 2.1$$

Question 3:

Find the domain of the function $f(x) = \frac{x^2 + 2x + 1}{x^2 - 8x + 12}$

Answer 3:

The given function is $f(x) = \frac{x^2 + 2x + 1}{x^2 - 8x + 12}$

$$f(x) = \frac{x^2 + 2x + 1}{x^2 - 8x + 12} = \frac{x^2 + 2x + 1}{(x - 6)(x - 2)}$$

It can be seen that function f is defined for all real numbers except at $x = 6$ and $x = 2$. Hence, the domain of f is $\mathbf{R} - \{2, 6\}$.

Question 4:


Find the domain and the range of the real function f defined by

$$f(x) = \sqrt{(x - 1)}$$

Answer 4:

The given real function is $f(x) = \sqrt{(x - 1)}$

It can be seen that $\sqrt{(x - 1)}$ is defined for $x \geq 1$.



Mathematics

(www.tiwariacademy.com : A step towards free education)

(Chapter – 2) (Relations and Functions)

(Class – XI)

Therefore, the domain of f is the set of all real numbers greater than or equal to 1 i.e., the domain of $f = [1, \infty)$.

$$\text{As } x \geq 1 \Rightarrow (x - 1) \geq 0 \Rightarrow \sqrt{(x - 1)} \geq 0$$

Therefore, the range of f is the set of all real numbers greater than or equal to 0 i.e., the range of $f = [0, \infty)$.

Question 5:

Find the domain and the range of the real function f defined by $f(x) = |x - 1|$.

Answer 5:

The given real function is $f(x) = |x - 1|$.

It is clear that $|x - 1|$ is defined for all real numbers.

\therefore Domain of $f = \mathbf{R}$

Also, for $x \in \mathbf{R}$, $|x - 1|$ assumes all real numbers.

Hence, the range of f is the set of all non-negative real numbers.

Question 6:

$$\text{Let } f = \left\{ \left(x, \frac{x^2}{1+x^2} \right) : x \in \mathbf{R} \right\}$$

be a function from \mathbf{R} into \mathbf{R} . Determine the range of f .

Answer 6:

$$\begin{aligned} f &= \left\{ \left(x, \frac{x^2}{1+x^2} \right) : x \in \mathbf{R} \right\} \\ &= \left\{ (0, 0), \left(\pm 0.5, \frac{1}{5} \right), \left(\pm 1, \frac{1}{2} \right), \left(\pm 1.5, \frac{9}{13} \right), \left(\pm 2, \frac{4}{5} \right), \left(3, \frac{9}{10} \right), \left(4, \frac{16}{17} \right), \dots \right\} \end{aligned}$$

The range of f is the set of all second elements. It can be observed that all these elements are greater than or equal to 0 but less than 1.

[Denominator is greater numerator] Thus, range of $f = [0, 1)$

Mathematics

(www.tiwariacademy.com : A step towards free education)

(Chapter – 2) (Relations and Functions)

(Class – XI)

Question 7:

Let $f, g: \mathbf{R} \rightarrow \mathbf{R}$ be defined, respectively by $f(x) = x + 1$, $g(x) = 2x - 3$. Find $f + g$, $f - g$ and $\frac{f}{g}$.

Answer 7:

$f, g: \mathbf{R} \rightarrow \mathbf{R}$ is defined as $f(x) = x + 1$, $g(x) = 2x - 3$

$$(f + g)(x) = f(x) + g(x) = (x + 1) + (2x - 3) = 3x - 2$$

$$\therefore (f + g)(x) = 3x - 2$$

$$(f - g)(x) = f(x) - g(x) = (x + 1) - (2x - 3) = x + 1 - 2x + 3 = -x + 4$$

$$\therefore (f - g)(x) = -x + 4$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}, g(x) \neq 0, x \in \mathbf{R}$$

$$\therefore \left(\frac{f}{g}\right)(x) = \frac{x+1}{2x-3}, 2x-3 \neq 0 \text{ or } 2x \neq 3$$

$$\therefore \left(\frac{f}{g}\right)(x) = \frac{x+1}{2x-3}, x \neq \frac{3}{2}$$

Mathematics

(www.tiwariacademy.com : A step towards free education)

(Chapter – 2) (Relations and Functions)

(Class – XI)

Question 8:

Let $f = \{(1, 1), (2, 3), (0, -1), (-1, -3)\}$ be a function from \mathbf{Z} to \mathbf{Z} defined by $f(x) = ax + b$, for some integers a, b . Determine a, b .

Answer 8:

$f = \{(1, 1), (2, 3), (0, -1), (-1, -3)\}$ and $f(x) = ax + b$

$$(1, 1) \in f \Rightarrow f(1) = 1 \Rightarrow a \times 1 + b = 1 \\ \Rightarrow a + b = 1$$

$$(0, -1) \in f \Rightarrow f(0) = -1 \Rightarrow a \times 0 + b = -1 \\ \Rightarrow b = -1$$

On substituting $b = -1$ in $a + b = 1$,

We obtain $a + (-1) = 1 \Rightarrow a = 1 + 1 = 2$. Thus, the respective values of a and b are 2 and -1.

Question 9:

Let R be a relation from \mathbf{N} to \mathbf{N} defined by $R = \{(a, b): a, b \in \mathbf{N} \text{ and } a = b^2\}$. Are the following true?

- (i) $(a, a) \in R$, for all $a \in \mathbf{N}$
 - (ii) $(a, b) \in R$, implies $(b, a) \in R$
 - (iii) $(a, b) \in R, (b, c) \in R$ implies $(a, c) \in R$.
- Justify your answer in each case.

Answer 9:

$R = \{(a, b): a, b \in \mathbf{N} \text{ and } a = b^2\}$

(i) It can be seen that $2 \in \mathbf{N}$; however, $2 \neq 2^2 = 4$.

Therefore, the statement " $(a, a) \in R$, for all $a \in \mathbf{N}$ " is not true.

(ii) It can be seen that $(9, 3) \in \mathbf{N}$ because $9, 3 \in \mathbf{N}$ and $9 = 3^2$. Now, $3 \neq 9^2 = 81$; therefore, $(3, 9) \notin \mathbf{N}$

Therefore, the statement " $(a, b) \in R$, implies $(b, a) \in R$ " is not true.

(iii) It can be seen that $(9, 3) \in R, (16, 4) \in R$ because $9, 3, 16, 4 \in \mathbf{N}$ and $9 = 3^2$ and $16 = 4^2$.

Now, $9 \neq 4^2 = 16$; therefore, $(9, 4) \notin \mathbf{N}$

Therefore, the statement " $(a, b) \in R, (b, c) \in R$ implies $(a, c) \in R$ " is not true.

Mathematics

(www.tiwariacademy.com : A step towards free education)

(Chapter – 2) (Relations and Functions)

(Class – XI)

Question 10:

Let $A = \{1, 2, 3, 4\}$, $B = \{1, 5, 9, 11, 15, 16\}$ and $f = \{(1, 5), (2, 9), (3, 1), (4, 5), (2, 11)\}$. Are the following true?

(i) f is a relation from A to B

(ii) f is a function from A to B .

Justify your answer in each case.

Answer 10:

$A = \{1, 2, 3, 4\}$ and $B = \{1, 5, 9, 11, 15, 16\}$

$\therefore A \times B = \{(1, 1), (1, 5), (1, 9), (1, 11), (1, 15), (1, 16), (2, 1), (2, 5), (2, 9), (2, 11), (2, 15), (2, 16), (3, 1), (3, 5), (3, 9), (3, 11), (3, 15), (3, 16), (4, 1), (4, 5), (4, 9), (4, 11), (4, 15), (4, 16)\}$

It is given that $f = \{(1, 5), (2, 9), (3, 1), (4, 5), (2, 11)\}$

(i) A relation from a non-empty set A to a non-empty set B is a subset of the Cartesian product $A \times B$.

It is observed that f is a subset of $A \times B$.

Thus, f is a relation from A to B .

(ii) Since the same first element i.e., 2 corresponds to two different images i.e., 9 and 11, relation f is not a function.

Question 11:

Let f be the subset of $\mathbf{Z} \times \mathbf{Z}$ defined by $f = \{(ab, a + b) : a, b \in \mathbf{Z}\}$. Is f a function from \mathbf{Z} to \mathbf{Z} : justify your answer.

Answer 11:

The relation f is defined as $f = \{(ab, a + b) : a, b \in \mathbf{Z}\}$

We know that a relation f from a set A to a set B is said to be a function if every element of set A has unique images in set B .

Since $2, 6, -2, -6 \in \mathbf{Z}$, $(2 \times 6, 2 + 6)$, $(-2 \times -6, -2 + (-6)) \in f$ i.e., $(12, 8)$, $(12, -8) \in f$

It can be seen that the same first element i.e., 12 corresponds to two different images i.e., 8 and -8. Thus, relation f is not a function.

Mathematics

(www.tiwariacademy.com : A step towards free education)

(Chapter – 2) (Relations and Functions)

(Class – XI)

Question 12:

Let $A = \{9, 10, 11, 12, 13\}$ and let $f: A \rightarrow \mathbf{N}$ be defined by $f(n) =$ the highest prime factor of n . Find the range of f .

Answer 12:

$A = \{9, 10, 11, 12, 13\}$ $f: A \rightarrow \mathbf{N}$ is defined as $f(n) =$ The highest prime factor of n

Prime factor of 9 = 3

Prime factors of 10 = 2, 5

Prime factor of 11 = 11

Prime factors of 12 = 2, 3

Prime factor of 13 = 13

$\therefore f(9) =$ The highest prime factor of 9 = 3

$f(10) =$ The highest prime factor of 10 = 5

$f(11) =$ The highest prime factor of 11 = 11

$f(12) =$ The highest prime factor of 12 = 3

$f(13) =$ The highest prime factor of 13 = 13

The range of f is the set of all $f(n)$, where $n \in A$.

\therefore Range of $f = \{3, 5, 11, 13\}$

