

Science

(www.tiwariacademy.net)

(Chapter 9)(Force and Laws of Motion)

Class - 9

Exercise

Question 1:

An object experiences a net zero external unbalanced force. Is it possible for the object to be travelling with a non-zero velocity? If yes, state the conditions that must be placed on the magnitude and direction of the velocity. If no, provide a reason.

Answer 1:

Yes. Even when an object experiences a net zero external unbalanced force, it is possible that the object is travelling with a non-zero velocity. This is possible only when the object has been moving with a constant velocity in a particular direction. Then, there is no net unbalanced force applied on the body. The object will keep moving with a non-zero velocity. To change the state of motion, a net non-zero external unbalanced force must be applied on the object.

Question 2:

When a carpet is beaten with a stick, dust comes out of it. Explain.

Answer 2:

Inertia of an object tends to resist any change in its state of rest or state of motion. When a carpet is beaten with a stick, then the carpet comes to motion. But, the dust particles try to resist their state of rest. According to Newton's first law of motion, the dust particles stay in a state of rest, while the carpet moves. Hence, the dust particles come out of the carpet.

Question 3:

Why is it advised to tie any luggage kept on the roof of a bus with a rope?

Answer 3:

When the bus accelerates and moves forward, it acquires a state of motion. However, the luggage kept on the roof, owing to its inertia, tends to remain in its state of rest. Hence, with the forward movement of the bus, the luggage tends to remain at its original position and ultimately falls from the roof of the bus. To avoid this, it is advised to tie any luggage kept on the roof of a bus with a rope.

www.tiwariacademy.com

A Free web support in education

Science

(www.tiwariacademy.net)

(Chapter 9)(Force and Laws of Motion)

Class - 9

Question 4:

A batsman hits a cricket ball which then rolls on a level ground. After covering a short distance, the ball comes to rest. The ball slows to a stop because

- (a) the batsman did not hit the ball hard enough.
- (b) velocity is proportional to the force exerted on the ball.
- (c) there is a force on the ball opposing the motion.
- (d) there is no unbalanced force on the ball, so the ball would want to come to rest.

Answer 4:

(c) A batsman hits a cricket ball, which then rolls on a level ground. After covering a short distance, the ball comes to rest because there is frictional force on the ball opposing its motion.

Frictional force always acts in the direction opposite to the direction of motion. Hence, this force is responsible for stopping the cricket ball.

Question 5:

A truck starts from rest and rolls down a hill with a constant acceleration. It travels a distance of 400 m in 20 s. Find its acceleration. Find the force acting on it if its mass is 7 metric tonnes (*Hint: 1 metric tonne = 1000 kg.*)

Answer 5:

Initial velocity of the truck, $u = 0$ m/s

Time taken, $t = 20$ s

Distance covered by the stone, $s = 400$ m

According to the second equation of motion:

$$s = ut + \frac{1}{2}at^2$$

We have

$$400 = 0 \times 20 + \frac{1}{2} \times a \times (20)^2$$

$$\Rightarrow 400 = 200a$$

$$\Rightarrow a = 2 \text{ m/s}^2$$

Now, Force = mass \times acceleration

$$\Rightarrow F = 7000 \times 2 = 14000 \text{ N}$$

www.tiwariacademy.com

A Free web support in education

Science

(www.tiwariacademy.net)
(Chapter 9)(Force and Laws of Motion)
Class - 9

Question 6:

A stone of 1 kg is thrown with a velocity of 20 m s^{-1} across the frozen surface of a lake and comes to rest after travelling a distance of 50 m. What is the force of friction between the stone and the ice?

Answer 6:

Initial velocity of the stone, $u = 20 \text{ m/s}$

Final velocity of the stone, $v = 0 \text{ m/s}$

Distance covered by the stone, $s = 50 \text{ m}$

According to the third equation of motion:

$$v^2 = u^2 + 2as$$

Where,

Acceleration, a

$$(0)^2 = (20)^2 + 2 \times a \times 50$$

$$a = -4 \text{ m/s}^2$$

The negative sign indicates that acceleration is acting against the motion of the stone.

Mass of the stone, $m = 1 \text{ kg}$

From Newton's second law of motion:

Force,

$$F = \text{Mass} \times \text{Acceleration}$$

$$F = ma$$

$$F = 1 \times (-4) = -4 \text{ N}$$

Hence, the force of friction between the stone and the ice is -4 N .

Science

(www.tiwariacademy.net)

(Chapter 9)(Force and Laws of Motion)

Class - 9

Question 7:

A 8000 kg engine pulls a train of 5 wagons, each of 2000 kg, along a horizontal track. If the engine exerts a force of 40000 N and the track offers a friction force of 5000 N, then calculate:

- (a) the net accelerating force;
- (b) the acceleration of the train; and
- (c) the force of wagon 1 on wagon 2.

Answer 7:

(a) 35000 N (b) 1.944 m/s^2 (c) 15552 N

(a) Force exerted by the engine, $F = 40000 \text{ N}$

Frictional force offered by the track, $F_f = 5000 \text{ N}$

Net accelerating force, $F_a = F - F_f = 40000 - 5000 = 35000 \text{ N}$

Hence, the net accelerating force is 35000 N.

(b) Acceleration of the train = a

The engine exerts a force of 40000 N on all the five wagons.

Net accelerating force on the wagons, $F_a = 35000 \text{ N}$

Mass of the wagons, $m = \text{Mass of a wagon} \times \text{Number of wagons}$

Mass of a wagon = 2000 kg

Number of wagons = 5

$\therefore m = 2000 \times 5 = 10000 \text{ kg}$

Total mass (including the mass of engine), $M = m + 8000 = 18000 \text{ kg}$

$F_a = Ma$

$$\Rightarrow a = \frac{F_a}{M} = \frac{35000}{18000} = 1.944 \text{ m/s}^2$$

(c) Mass of 4 wagons (excluding wagon 1) $m = 2000 \times 4 = 8000 \text{ kg}$

Acceleration of the train = 1.944 m/s^2

\therefore The force of wagon 1 on wagon 2 = $ma = 8000 \times 1.944 = 15552 \text{ N}$

www.tiwariacademy.com

A Free web support in education

Science

(www.tiwariacademy.net)

(Chapter 9)(Force and Laws of Motion)

Class - 9

Question 8:

An automobile vehicle has a mass of 1500 kg. What must be the force between the vehicle and road if the vehicle is to be stopped with a negative acceleration of 1.7 ms^{-2} ?

Answer 8:

Mass of the automobile vehicle, $m = 1500 \text{ kg}$

Final velocity, $v = 0 \text{ m/s}$

Acceleration of the automobile, $a = -1.7 \text{ ms}^{-2}$

From Newton's second law of motion:

Force = Mass \times Acceleration = $1500 \times (-1.7) = -2550 \text{ N}$

Hence, the force between the automobile and the road is -2550 N , in the direction opposite to the motion of the automobile.

Question 9:

What is the momentum of an object of mass m , moving with a velocity v ?

(a) $(mv)^2$ (b) mv^2 (c) $\frac{1}{2}mv^2$ (d) mv

Answer 9:

(d) mv

Mass of the object = m

Velocity = v

Momentum = Mass \times Velocity

Momentum = mv

Question 10:

Using a horizontal force of 200 N, we intend to move a wooden cabinet across a floor at a constant velocity. What is the friction force that will be exerted on the cabinet?

Answer 10:

A force of 200 N is applied in the forward direction. Thus, from Newton's third law of motion, an equal amount of force will act in the opposite direction. This opposite force is the frictional force exerted on the cabinet. Hence, a frictional force of 200 N is exerted on the cabinet.

www.tiwariacademy.com

A Free web support in education

Science

(www.tiwariacademy.net)

(Chapter 9)(Force and Laws of Motion)

Class - 9

Question 11:

Two objects, each of mass 1.5 kg are moving in the same straight line but in opposite directions. The velocity of each object is 2.5 ms^{-1} before the collision during which they stick together. What will be the velocity of the combined object after collision?

Answer 11:

Mass of one of the objects, $m_1 = 1.5 \text{ kg}$

Mass of the other object, $m_2 = 1.5 \text{ kg}$

Velocity of m_1 before collision, $v_1 = 2.5 \text{ m/s}$

Velocity of m_2 , moving in opposite direction before collision, $v_2 = -2.5 \text{ m/s}$

(Negative sign arises because mass m_2 is moving in an opposite direction)

After collision, the two objects stick together.

Total mass of the combined object = $m_1 + m_2$

Velocity of the combined object = v

According to the law of conservation of momentum:

Total momentum before collision = Total momentum after collision

$$m_1 v_1 + m_2 v_2 = (m_1 + m_2) v$$

$$1.5(2.5) + 1.5(-2.5) = (1.5 + 1.5) v$$

$$3.75 - 3.75 = 3 v$$

$$v = 0$$

Hence, the velocity of the combined object after collision is 0 m/s.

www.tiwariacademy.com

A Free web support in education

Science

(www.tiwariacademy.net)

(Chapter 9)(Force and Laws of Motion)

Class - 9

Question 12:

According to the third law of motion when we push on an object, the object pushes back on us with an equal and opposite force. If the object is a massive truck parked along the roadside, it will probably not move. A student justifies this by answering that the two opposite and equal forces cancel each other. Comment on this logic and explain why the truck does not move.

Answer 12:

The truck has a large mass. Therefore, the static friction between the truck and the road is also very high. To move the car, one has to apply a force more than the static friction. Therefore, when someone pushes the truck and the truck does not move, then it can be said that the applied force in one direction is cancelled out by the frictional force of equal amount acting in the opposite direction.

Therefore, the student is right in justifying that the two opposite and equal cancel each other.

Question 13:

A hockey ball of mass 200 g travelling at 10 m s^{-1} is struck by a hockey stick so as to return it along its original path with a velocity at 5 m s^{-1} . Calculate the change of momentum occurred in the motion of the hockey ball by the force applied by the hockey stick.

Answer 13:

Mass of the hockey ball, $m = 200 \text{ g} = 0.2 \text{ kg}$

Hockey ball travels with velocity, $v_1 = 10 \text{ m/s}$

Initial momentum = mv_1

Hockey ball travels in the opposite direction with velocity, $v_2 = -5 \text{ m/s}$

Final momentum = mv_2

Change in momentum = $mv_1 - mv_2 = 0.2 [10 - (-5)] = 0.2 (15) = 3 \text{ kg m s}^{-1}$

Hence, the change in momentum of the hockey ball is 3 kg m s^{-1} .

www.tiwariacademy.com

A Free web support in education

Science

(www.tiwariacademy.net)

(Chapter 9)(Force and Laws of Motion)

Class - 9

Question 14:

A bullet of mass 10 g travelling horizontally with a velocity of 150 ms^{-1} strikes a stationary wooden block and comes to rest in 0.03 s. Calculate the distance of penetration of the bullet into the block. Also calculate the magnitude of the force exerted by the wooden block on the bullet.

Answer 14:

Now, it is given that the bullet is travelling with a velocity of 150 m/s.

Thus, when the bullet enters the block, its velocity = Initial velocity, $u = 150 \text{ m/s}$

Final velocity, $v = 0$ (since the bullet finally comes to rest)

Time taken to come to rest, $t = 0.03 \text{ s}$

According to the first equation of motion, $v = u + at$

Acceleration of the bullet, a

$$0 = 150 + (a \times 0.03 \text{ s})$$

$$a = \frac{-150}{0.03} = -5000 \text{ m/s}^2$$

(Negative sign indicates that the velocity of the bullet is decreasing.)

According to the third equation of motion:

$$v^2 = u^2 + 2as$$

$$0 = (150)^2 + 2(-5000)s$$

$$s = \frac{-(150)^2}{-2(5000)} = \frac{22500}{10000} = 2.25 \text{ m}$$

Hence, the distance of penetration of the bullet into the block is 2.25 m.

From Newton's second law of motion:

Force, $F = \text{Mass} \times \text{Acceleration}$

Mass of the bullet, $m = 10 \text{ g} = 0.01 \text{ kg}$

Acceleration of the bullet, $a = 5000 \text{ m/s}^2$

$$F = ma = 0.01 \times 5000 = 50 \text{ N}$$

www.tiwariacademy.com

A Free web support in education

Science

(www.tiwariacademy.net)
(Chapter 9)(Force and Laws of Motion)
Class - 9

Question 15:

An object of mass 1 kg travelling in a straight line with a velocity of 10 ms^{-1} collides with, and sticks to, a stationary wooden block of mass 5 kg. Then they both move off together in the same straight line. Calculate the total momentum just before the impact and just after the impact. Also, calculate the velocity of the combined object.

Answer 15:

Mass of the object, $m_1 = 1 \text{ kg}$

Velocity of the object before collision, $v_1 = 10 \text{ m/s}$

Mass of the stationary wooden block, $m_2 = 5 \text{ kg}$

Velocity of the wooden block before collision, $v_2 = 0 \text{ m/s}$

\therefore Total momentum before collision = $m_1 v_1 + m_2 v_2$

$$= 1 (10) + 5 (0) = 10 \text{ kg m s}^{-1}$$

It is given that after collision, the object and the wooden block stick together.

Total mass of the combined system = $m_1 + m_2$

Velocity of the combined object = v

According to the law of conservation of momentum:

Total momentum before collision = Total momentum after collision

$$m_1 v_1 + m_2 v_2 = (m_1 + m_2) v$$

$$1 (10) + 5 (0) = (1 + 5) v$$

$$v = \frac{10}{6} = \frac{5}{3} \text{ m/s}$$

The total momentum after collision = 10 and velocity of combined object is $\frac{5}{3} \text{ m/s}$.

Science

(www.tiwariacademy.net)

(Chapter 9)(Force and Laws of Motion)

Class - 9

Question 16:

An object of mass 100 kg is accelerated uniformly from a velocity of 5 ms^{-1} to 8 ms^{-1} in 6 s. Calculate the initial and final momentum of the object. Also, find the magnitude of the force exerted on the object.

Answer 16:

Initial velocity of the object, $u = 5 \text{ m/s}$

Final velocity of the object, $v = 8 \text{ m/s}$

Mass of the object, $m = 100 \text{ kg}$

Time take by the object to accelerate, $t = 6 \text{ s}$

Initial momentum = $mu = 100 \times 5 = 500 \text{ kg ms}^{-1}$

Final momentum = $mv = 100 \times 8 = 800 \text{ kg ms}^{-1}$

Force exerted on the object, $F = \frac{mv - mu}{t}$

$$= \frac{m(v - u)}{t} = \frac{800 - 500}{6} = \frac{300}{6} = 50 \text{ N}$$

Initial momentum of the object is 500 kg ms^{-1} .

Final momentum of the object is 800 kg ms^{-1} .

Force exerted on the object is 50 N.

Question 17:

Akhtar, Kiran and Rahul were riding in a motorocar that was moving with a high velocity on an expressway when an insect hit the windshield and got stuck on the windscreen. Akhtar and Kiran started pondering over the situation. Kiran suggested that the insect suffered a greater change in momentum as compared to the change in momentum of the motorocar (because the change in the velocity of the insect was much more than that of the motorocar). Akhtar said that since the motorocar was moving with a larger velocity, it exerted a larger force on the insect. And as a result the insect died. Rahul while putting an entirely new explanation said that both the motorocar and the insect experienced the same force and a change in their momentum. Comment on these suggestions.

www.tiwariacademy.com

A Free web support in education

Science

(www.tiwariacademy.net)

(Chapter 9)(Force and Laws of Motion)

Class - 9

Answer 17:

According to the law of conservation of momentum:

Momentum of the car and insect system before collision = Momentum of the car and insect system after collision

Hence, the change in momentum of the car and insect system is zero.

The insect gets stuck on the windscreen. This means that the direction of the insect is reversed. As a result, the velocity of the insect changes to a great amount. On the other hand, the car continues moving with a constant velocity. Hence, Kiran's suggestion that the insect suffers a greater change in momentum as compared to the car is correct. The momentum of the insect after collision becomes very high because the car is moving at a high speed. Therefore, the momentum gained by the insect is equal to the momentum lost by the car.

Akhtar made a correct conclusion because the mass of the car is very large as compared to the mass of the insect.

Rahul gave a correct explanation as both the car and the insect experienced equal forces caused by the Newton's action-reaction law. But, he made an incorrect statement as the system suffers a change in momentum because the momentum before the collision is equal to the momentum after the collision.

Question 18:

How much momentum will a dumbbell of mass 10 kg transfer to the floor if it falls from a height of 80 cm? Take its downward acceleration to be 10 m s^{-2} .

Answer 18:

Mass of the dumbbell, $m = 10 \text{ kg}$

Distance covered by the dumbbell, $s = 80 \text{ cm} = 0.8 \text{ m}$

Acceleration in the downward direction, $a = 10 \text{ m/s}^2$

Initial velocity of the dumbbell, $u = 0$

Final velocity of the dumbbell (when it was about to hit the floor) = v

www.tiwariacademy.com

A Free web support in education

Science

(www.tiwariacademy.net)

(Chapter 9)(Force and Laws of Motion)

Class - 9

According to the third equation of motion:

$$v^2 = u^2 + 2as$$

$$v^2 = 0 + 2 (10) 0.8$$

$$v = 4 \text{ m/s}$$

Hence, the momentum with which the dumbbell hits the floor is
 $= mv$

$$= 10 \times 4 \text{ kg m s}^{-1}$$

$$= 40 \text{ kg m s}^{-1}$$